

Description of the final instar larva of *Calopteryx exul* Sélys, 1853 (Zygoptera: Calopterygidae)

Rassim Khelifa*

Département d'Écologie et du Génie de l'Environnement, Faculté des Sciences de la Nature et de la Vie et
des Sciences de la Terre et de l'Univers, Université 08 Mai 1945, Guelma 24000, Algeria

(Received 12 March 2012; final version received 27 April 2012)

The last instar larva of *Calopteryx exul* Sélys is described and illustrated based on larvae collected from the Seybouse River (northeast Algeria) and reared in the laboratory. A comparative analysis of three other congeneric species is presented.

Keywords: Odonata; damselflies; Calopterygidae; endemic species; endangered; exuvia; taxonomy

Introduction

Calopteryx exul Sélys, 1853 is one of the most threatened zygopterans in the Mediterranean basin with a geographic distribution restricted to North Africa (Tunisia, Algeria, and Morocco) (Riservato et al., 2009). In Algeria, the species has not been found since Martin (1910), but recently a colony was discovered in the Seybouse River (northeast Algeria) (Khelifa et al., 2011). This represents the only population currently known in the country and the largest in the Maghreb (Khelifa, in press). The adult preferred habitat is large watercourses with relatively shallow and fast flowing water (Khelifa, in press).

To date, no description of the larvae has been published except for a general illustration (sketch) by Jödicke et al. (2000) of one exuvia collected in northern Tunisia, stating that the species is characterized by rather narrow forewing sheaths with caudal appendages rather long compared to the length of antennae or labium. Here, I present a detailed description of *C. exul* exuvia based on individuals collected from northeast Algeria.

Methods

Study area

The study was conducted in the Seybouse River, northeast Algeria. This river is formed by the junction of Cherf and Bouhamdane wadi at Medjez Amar (36°26'35" N, 7°18'39" E) and flows

*Email: rassimkhelifa@gmail.com

51 into the Mediterranean at Sidi Salem (36°52'3" N, 7°46'25" E) near Annaba. The hydroperiod is
 52 similar to that of Numidia with a wet season between October and May and a dry season between
 53 June and September. The annual rainfall varies from 450 mm upstream to 735 mm downstream
 54 (ABHCSM, 2009). The river edges are usually covered by *Typha angustifolia*, *Tamarix gallica*,
 55 *Nerium oleander*, and *Rubus ulmifolius*.

57 **Larval sampling and rearing**

58 I used previous information on adult distribution of *Calopteryx* species in the Seybouse watershed
 59 to guide collection of larvae (Khelifa et al., 2011). Forty-one larvae were collected in April 2010
 60 and 2011 with a hand net of 0.5 mm mesh from 10 m stretches of the shore in three sites (Table 1).
 61 I chose only large larvae to reduce mortality risks during rearing. Samples were brought to the
 62 laboratory within 2 hours of collection.

63 In the laboratory, I followed Van Gossum et al. (2003) to increase rearing success. Larvae were
 64 reared individually in oxygenated water and kept at room temperature of $18 \pm 3^\circ\text{C}$ under natural
 65 light conditions. We submerged a few sticks in water in order to help larvae to climb and emerge.
 66 Aquaria were covered with a net so that teneral could not fly away after emergence. A large
 67 range of prey was provided, namely daphniids, dipteran and mayfly larvae, amphipods, and small
 68 tadpoles. I checked the aquaria daily for the presence of exuviae and teneral. To limit damage to
 69 the population, all the emerged adults were reintroduced to their original localities within 24 h of
 70 emergence.

71 To calculate ratios, I measured the scape and pedicel length, the distance between antennomere
 72 2 and antennomere 7, the length of the prementum, posterior tibia, and posterior femur using the
 73 same limits as Heidemann & Seidenbusch (2002). Values are presented as mean \pm SD.

77 **Results**

78 **Habitat**

79 Sites where *Calopteryx exul* larvae were collected were wide stretches with relatively shallow
 80 and fast flowing water (Table 1). Larvae were abundant at the edge where vegetation, dominated
 81 by *Schoenoplectus maritimus* and *Typha angustifolia*, was hanging over the bank. Other species
 82 of odonates found included *Gomphus lucasii*, *Onychogomphus costae*, *Platycnemis subdilatata*,
 83 and rarely *Calopteryx haemorrhoidalis*.

84 **Description** (Figures 1–8)

85 All larvae emerged between 23:00 and 04:00 except for one individual which emerged at 19:30.
 86 From 41 larvae collected only 32 emerged successfully. All measurements and ratios below are
 87 based on the final instar exuviae of these emerged individuals.

88 **Body.** Yellowish to brownish, or rarely darker, slender body, mean total length (excluding
 89 antennae and gills) of 21.61 ± 1.35 mm (Figure 1a). Almost totally covered by small setae
 90 (Figure 1b).

91 **Head.** Wider than long (mean width of 4.67 ± 0.33 mm) and almost totally covered by sparse
 92 patches of small setae. Vertex with three conspicuous ocelli. Prominent postocular projections
 93 (occipital tubercles) approximately hemispheric (bulge-like) in lateral view (Figure 2a) but blunt
 94




Figure 1. Body of *Calopteryx exul* final instar exuvia: (a) dorsal view; (b) enlarged dorsal view of abdominal segment 4 showing setae of body surface.

in dorsal view (Figure 2b). Antennal sclerites large, surrounded by rather long setae. Antennae 7-segmented. Scape (first antennomere) exceeding the length of all remaining antennomeres (ratio 1.5:1), triqueteral (3 edged) except distal part, which is cylindrical. Segments 2 to 6 cylindrical, segment 7 conical. Each scape with longitudinal mediodorsal pale stripe; Scape and pedicel (second antennomere) covered with furry layer of short setae. No gap between insertion of labial palpus to prementum. Two palpal setae close to insertion of long movable hook (Figure 3). Prementum deeply divided by long median cleft and with pair of short premental setae ventrally disposed in its distal part (median lobes) (Figure 4).

Thorax: Mediodistal corners of pronotum relatively blunt. Wing cases with approximately semi-circular tips. Ratio of hind wing case length: width 3.6:1. Forewing cases reaching anterior margin of S4, hind wing cases exceed it and rarely reach its posterior margin.

Legs. Long with two dark bands on femur and tibia. Tibia always slightly longer than femur (ratio of posterior tibia:posterior femur 1.1:1). Tarsi 3 segmented, with dense, relatively long setae on first segment and 2 ventral parallel lines of setae on segments 2 and 3 (Figure 7).

151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200




Figure 2. Head of *Calopteryx exul*: (a) lateral view; (b) dorsal view.






Figure 3. Left labial palpus. Because the intermediate hook crosses the movable hook, a medioventral view is shown so that all hooks are visible.

201
202
203
204
205
206
207
208
209
210
211
212
213
214




215 Figure 4. Prementum, view of inner surface.

216
217
218
219
220
221
222
223
224
225
226
227
228
229



230 Figure 5. Male valves, ventral view.

231
232
233
234
235
236
237
238
239
240
241
242
243



244 Figure 6. Female ovipositor and gonapophyses: (a) lateral view; (b) ventral view.

245
246
247
248
249
250

Abdomen. Lateral flanges clearly evident. Male primary genitalia (valves) straight, slender, not exceeding posterior margin of 9th abdominal segment (Figure 5). In females, ovipositor not reaching posterior margin of final abdominal segment, both extremities of lateral gonapophyses in contact (Figure 6a, b).

251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300




Figure 7. Left metatarsus, oblique ventral view.




Figure 8. Caudal gills: (a) right lateral gill; a small part of the gill edge is enlarged showing the curved spines; (b) median gill.

Caudal gills. Triquetral lateral gills and lamellate median gill covered with short setae. Lateral gills always longer than the median (ratio 1.5:1). The three edges of lateral gills and the ventral and dorsal margins of the median gill with tiny, equidistant, curved spines (Figure 8a). Both lateral and median gill with blunt apex and are marked with one pale spot (sometimes two spots on lateral gill) usually in the middle of each gill (Figure 8a, b); second pale spot of lateral gill, if present, situated between first spot and the apex, obvious in larva but usually fainter or totally absent in final instar exuvia.

Discussion

In this paper I describe the last instar exuviae of the endangered, river-dwelling, *Calopteryx exul*. In Algeria, like in the rest of its geographic range, the species has undergone great habitat loss (Riservato et al., 2009) and the lack of data about its life history, ecology, and larval taxonomy has made it difficult to establish a regional conservation plan.

A comparative analysis of *C. exul* with three other Palearctic species described by Heidemann and Seidenbusch (2002), namely *C. splendens* (Harris, 1782), *C. haemorrhoidalis* (Vander Linden, 1825), and *C. virgo* (Linnaeus, 1758) is presented in Table 2. *C. exul* exuviae are most similar to *C. splendens* in a set of traits and ratios. They have the same body size and similar ratios of mentum:tibia, scape:pedicel, and scape:remaining antennomeres (Table 2). Also, females have a similar shape and disposition of the ovipositor and gonapophyses (Heidemann & Seidenbusch, 2002). One clear difference between these two species is that the male valves are short, thick, and have hemispheric extremities in *C. splendens*. Further taxonomic efforts are necessary to provide additional identification characters between these closely related species (Weekers et al., 2001). The distinction of both species should pose few taxonomic problems, and additionally they do not overlap in their geographic range; *C. exul* is endemic to the Maghreb whereas *C. splendens* is

301 widely distributed over Europe and Western Asia. Two other *splendens*-like taxa, *C. waterstoni*
 302 (Schneider, 1984) and *C. hyalina* (Martin, 1909), have fully hyaline wings similar to those of
 303 *C. exul*. According to recent data from Syria and Lebanon, *C. hyalina* is currently almost extinct
 304 (Riservato et al., 2009). Q6

305 Regarding *C. virgo*, besides its larger body size, almost all ratios were different. Among the
 306 four species presented in Table 2, *C. virgo* is particularly characterized by conspicuous conical
 307 and acute postocular tubercles (Heidemann & Seidenbusch, 2002) and thus could not be confused
 308 with the three other species.

309 However, particular attention should be paid to differentiate *C. exul* from *C. haemorrhoidalis*
 310 because these species coexist and emerge at the same sites (personal observation). According to
 311 Heidemann and Seidenbusch (2002), female ovipositor and male valves are the most important
 312 traits that can be used in the identification of *Calopteryx* species. *Calopteryx exul* and *C. haem-*
 313 *orrhoidalis* differ substantially in these two characters. Besides the fact that *C. exul* is longer and
 314 paler, the *C. haemorrhoidalis* ovipositor usually exceeds the final abdominal segment and the
 315 extremities of lateral gonapophyses do not touch (Heidemann & Seidenbusch, 2002). *Calopteryx*
 316 *haemorrhoidalis* males have rather acute slender valves that extend beyond the 9th abdominal
 317 segment (Heidemann & Seidenbusch, 2002). Using these traits might not be enough to accurately
 318 recognize the species in case of inter-population polymorphism, but adding the ratios in Table 2
 319 and some characters illustrated and described above (see results) will yield reliable identification.
 320 However, future studies using larger sample sizes could lead to new measures and ratios slightly
 321 different from those reported by Heidemann and Seidenbusch (2002) for the three species of
 322 *Calopteryx*.

323 Furthermore, setae of tarsi of *C. exul* are similar to those of *C. haemorrhoidalis* (personal
 324 observation) and *C. maculata* (M. May, personal communication). Since neither species is closely
 325 related to *C. exul* (Weekers et al., 2001), this is probably a general pattern in *Calopteryx*. Another
 326 trait which could be an important identification key is the gill spines. Angle of curvature, length,
 327 and disposition of gill spines are most likely to vary between species and might be taken into
 328 account for future description of *Calopteryx* species.

329 It is well known that using the last instar exuviae in odonatological studies gives important
 330 information on species life history, ecology (Corbet, 1999), and genetics (Watts, 2009). Moreover, Q7
 331 its occurrence at a given water body is a better indicator of reproduction than that of adults or
 332 larvae (Hardersen, 2008; Ott et al., 2007; Reabel et al., 2010; Samways et al., 2010).

333

334

335 Acknowledgements

336

337 I thank Pr. H. Dumont and Dr. G. Carchini for their useful comments and critics. I also thank Youcefi Abdeldjalil, Zebba Q8
 338 Rabah, Kahalerras Amin, and Amari Hichem for field assistance. Thanks to Pr. Boudjéma Samraoui for sharing some
 339 advice in larvae rearing. I am grateful to Pr. Jean-Pierre Boudot and Sónia Ferreira for their help in the documentation.

340

341

342 References

343

344 ABHCSM (Agence de Bassin Hydrographique Constantinois-Seybouse-Mellegue). (2009). Qualité des eaux superficielles
 345 dans les bassins du Kebir-Rhumel, de la Seybouse et de la Medjerda-Mellegue, 2004–2007 (N° 12). Q10

346

347 Corbet, P.S. (1999). *Dragonflies: Behaviour and Ecology of Odonata*. Colchester: Harley Books.

348

349 Hardersen, S. (2008). Dragonfly (Odonata) communities at three lotic sites with different hydrological characteristics.
 350 *Italian Journal of Zoology*, 75, 271–283.

351

352 Heidemann, H. & Seidenbusch, R. (2002). *Larves et exuvies des libellules de France et d'Allemagne (sauf de Corse)*. Q11
 353 Société Française d'Odonatologie.

354

355 Jödicke, R., Arlt, J., Kunz, B., Lopau, W., & Seidenbusch, R. (2000). The Odonata of Tunisia. *International Journal of*
 356 *Odonatology*, 3, 41–71.

357

358 Khelifa, R. (in press). Flight period, apparent sex ratio and habitat preferences of the Maghribian endemic *Calopteryx*
 359 *exul* Selys, 1853 (Odonata: Zygoptera). *Revue d'Ecologie (La Terre et la Vie)*. Q12

351 Khelifa, R., Youcefi, A., Kahlerras, A., Alfarhan, A., Al-Rasheid, K.A.S., & Samraoui, B. (2011). L'odonatofaune (Insecta
352 : Odonata) du bassin de la Seybouse en Algérie : intérêt pour la biodiversité du Maghreb. *Revue d'Ecologie (La Terre
353 et la Vie)*, 66, 55–66.

354 Ott, J., Schorr, M., Trockur, B., & Lingenfelder, U. (2007). Artenschutzprogramm Fur Die Gekielte Smaragdlibelle
355 (*Oxygastra Curtisii*) in Deutschland an Der Our/Species Protection Programme for the Orange-spotted Emerald
356 (*Oxygastra Curtisii*) in Germany. The Example of the Our River Population. *Invertebrate Ecology and Conservation
357 Monographs* (Vol. 3). Pensoft Publishers.

358 Raebel, E.M., Merckx, T., Riordan, P., Macdonald, D.W., & Thompson, D.J. (2010). The dragonfly delusion: why it is
359 essential to sample exuviae to avoid biased surveys. *Journal of Insect Conservation*, 14, 523–533.

360 Riservato, E., Boudot, J.-P., Ferreira, S., Jović, M., Kalkman, V.J., Schneider, W., Samraoui, B., & Cuttelod, A. (2009). The
361 status and distribution of dragonflies of the Mediterranean basin Gland, IUCN (International Union for Conservation
362 of Nature), Switzerland and Malaga, Spain: IUCN vii + 33 pp.

363 Samways, M.J., McGeoch, M.A., & New, T.R. (2010). *Insect conservation: Handbook of approaches and methods*.
364 Oxford: Oxford University Press.

365 Van Gossom, H., Sánchez, R., & Cordero-Rivera, A. (2003). Observations on rearing damselflies under laboratory
366 conditions. *Animal Biology*, 53, 37–45.

367 Watts, P.C., Thompson, D.J., Daguët, C., & Kemp, S.J. (2005). Exuviae as a reliable source of DNA for population-genetic
368 analysis of odonates. *Odonatologica*, 34, 183–187.

369 Weekers, P.H.H., De Jonckheere, J.F., & Dumont, H.J. (2001). Phylogenetic relationships inferred from ribosomal ITS
370 sequences *Calopteryx* (Insecta: Odonata) of the West Mediterranean and Adjacent West European Zone. *Molecular
371 Phylogenetics and Evolution*, 20, 89–99.

Q13

Q9

351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400